ΒΡΟΓΧΟΣΚΟΠΗΣΗ

Φίλιππος Εμμανουήλ

- Πνευμονολόγος Mediterraneo hospital Γλυφάδα
- Επιστημονικός συνεργάτης Μονάδας θωρακικής ενδοσκόπησης

Α’ Πνευμονολογικής Κλινικής ΕΚΠΑ



- Προετοιμασία για βρογχοσκόπηση – οδηγίες BTS
- Τοπική αναισθησία και νευροληπτοαναλγησία
- Ανατομία λάρυγγα, τραχείας και βρογχικού δένδρου- ανατομικές παραλλαγές και διαταραχές
- Ασφαλής διαγνωστική προσπέλαση ενδοβρογχικών βλαβών
- Αιμόπτυση και αντιμετώπιση
- Βρογχοσκόπηση για λοιμώξεις σε ανοσοεπαρκείς και ανοσοκατεσταλμένους.
- Ο ρόλος του βρογχοκυψελιδικού εκπλύματος (BAL)
- Σταδιοποίηση κατά TNM, οδηγίες ACCP-ESTS.
- Προσπέλαση λεμφαδενικών σταθμών με TBNA (ιστολογική βελόνα 19G - CT guided approach).
- Προσπέλαση λεμφαδενικών σταθμών με EBUS-TBNA. Συστηματικό assessment (EBUS, b-EUS).
- Διαβρογχική προσπέλαση περιφερικών όγκων με radial EBUS και ακτινοσκόπηση.
- Νεότερα δεδομένα για την πλοηγούμενη βρογχοσκόπηση

Question I.6 Originally, the flexible bronchoscope was designed to be held as shown in the Figure below. A reason for this is

A. The operator must always stand behind the patient; therefore it is best for the control section to be held in the left hand.
B. Dr. Ikeda, original designer of the flexible bronchoscope, was left-handed.
C. The operator must always stand to the right of the patient; therefore, it is best for the control section to be held in the left hand so that the bronchoscopist's right hand can be closest to the patient.
D. The operator must always stand to the left of the patient; therefore, it is best for the control section to be held in the left hand so that the bronchoscopist's right hand can be closest to the patient.

http://www.bronchoscopy.org/education

Προετοιμασία για βρογχοσκόπηση- οδηγίες BTS

Practitioners undertaking FB should be familiar with, and adhere to the national and local guidance for obtaining informed consent.

All patients undergoing bronchoscopy should have heart rate, respiratory rate, blood pressure and oxygen saturation recorded repeatedly, including before, during and after the procedure. (Grade D)

Continuous ECG monitoring should be used when there is a high clinical risk of arrhythmia. (Grade D)

Oxygen supplementation should be used when desaturation is significant (pulse oximeter oxygen saturation (SpO2)>4% change, or SpO2<90%) and prolonged (>1 min) to reduce the risk of hypoxaemia-related complications. (Grade D)

Du Rand IA, et al. Thorax 2013;68:i1–i44

Perform coagulation studies, platelet count and haemoglobin concentration when there are clinical risk factors for abnormal coagulation. (Grade D)

Bronchoscopy with lavage can be performed with platelet counts >20 000 per µL.

Stop warfarin 5 days prior to bronchoscopy.

Continue with LMWH.

Check if INR <1.5 prior.



Du Rand IA, et al. Thorax 2013;68:i1–i44

Discontinue clopidogrel 7 days prior to consideration of EBB and TBLB. Low-dose aspirin alone can be continued. (Grade C)



Du Rand IA, et al. Thorax 2013;68:i1–i44

ΑΣΦΑΛΕΙΑ ΤΗΣ ΒΡΟΓΧΟΣΚΟΠΗΣΗΣ ΣΕ ΕΙΔΙΚΟΥΣ ΠΛΗΘΥΣΜΟΥΣ

Asthma control should be optimised prior to bronchoscopy, especially when bronchoalveolar lavage (BAL) is likely to be performed. (Grade C)

Nebulised bronchodilators should be considered before bronchoscopy in patients with asthma.

Chronic obstructive pulmonary disease (COPD) treatment should be optimised prior to bronchoscopy. (Grade D)
Bronchoscopists should be cautious when sedating patients with COPD. (Grade D)

Ischaemic heart disease: Liaison with cardiologists should be considered in high-risk patients with cardiac disease.

FB should ideally be delayed for 4 weeks after MI. (Grade D)



Age alone should not be a contraindication for bronchoscopy. (Grade D)

Du Rand IA, et al. Thorax 2013;68:i1–i44

Question I.11: Which of the following topical anesthetics should be used in a patient who has just informed you of a severe allergic reaction to Novocaine during a recent visit to the dentist.

A. Lidocaine
B. Benzocaine
C. Tetracaine
D. Cocaine
E. All of the above

ΤΟΠΙΚΗ ΑΝΑΙΣΘΗΣΙΑ

Lidocaine should be used for topical anaesthesia during bronchoscopy, unless contraindicated. (Grade A)

Nasal topical anaesthesia is most effectively provided using 2% lidocaine gel. (Grade A)

Cricothyroid and spray-as-you-go techniques are effective in delivering lidocaine to the vocal cords and trachea. (Grade B)

Nebulisation is not recommended as a technique for delivering lidocaine to the airways. (Grade B)

1% lidocaine solution for spray-as-you-go administration. (Grade A)

To reduce the risk of lidocaine toxicity, use the lowest dose of lidocaine sufficient to prevent excessive coughing and provide patient comfort. (Grade D)

Remain vigilant for objective and subjective symptoms of lidocaine toxicity (Grade B) - monitor and document the total lidocaine dose delivered at all sites during bronchoscopy.

Du Rand IA, et al. Thorax 2013;68:i1–i44

Randomised studies suggesting that a low cough score is achievable with a total lidocaine dose of less than 160 mg.

Bronchoscopists should use the lowest dose of lidocaine possible, while still ensuring good bronchoscopic conditions and patient comfort.

Lidocaine 1% = 10 mg/mL (Percentage is measured in grams per 100 mL)

Δοσοεξαρτόμενες παρενέργειες: Αν ο ασθενής λάβει πάνω από 7-8 mg/kg
(Δεν ξεπερνάμε τα 50 ml ξυλοκαΐνης 1% σε ασθενή 70kg).

Du Rand IA, et al. Thorax 2013;68:i1–i44

ΤΟΞΙΚΟΤΗΤΑ ΑΠΌ ΛΙΔΟΚΑΙΝΗ – Κ.Ν.Σ και καρδιαγγειακό:

- Μεταλλική γεύση
- Ναυτία
- Οπτικές και ακουστικές διαταραχές- εμβοές
- Αποπροσανατολισμός
- Με μεγαλύτερες δόσεις- Καταστολή ΚΝΣ
- Σπασμοί μυών
- Επιληπτική κρίση
- Κώμα
- Αναπνευστική καταστολή
- Υπόταση
- Αρρυθμίες

- Μεθαιμοσφαιριναιμία – (κυανωση, δύσπνοια).

- **Αλλεργική αντίδραση**

Brown Emergency Medicine

American Society of Regional Anesthesia and Pain Medicine (ASRA)
Reg Anesth Pain Med. 2018; 43(2):150-153

ΑΝΤΙΜΕΤΩΠΙΣΗ ΤΟΞΙΚΟΤΗΤΑΣ ΑΠΌ ΛΙΔΟΚΑΙΝΗ

- Βενζοδιαζεπίνες (σπασμοί)
- Ενδοφλέβια υγρά και αγγειοσυσπαστικά
- Κατά προτίμηση επινεφρίνη bolus (≤1 mcg/kg).

- Intravenous (IV) infusion of lipid emulsions to reverse the cardiac and neurologic effects of local-anesthetic toxicity (Intralipid™ 20% 1.5 mL/kg (lean body weight) IV bolus over 2 – 3 minutes, followed by an infusion of Intralipid™ 20% 15 mL/kg IV over 60 minutes)

- HCO3 για σοβαρή οξέωση.

- Επίπεδα μεθαιμοαιμοσφαιρίνης στο αίμα. Για συμπτωματικούς ασθενείς ή ασθενείς με επίπεδα >25%, IV methylene blue (1 mg/kg σε 5 λεπτά). Η ανταπόριση είναι σύνήθως γρήγορη (<1 ώρα).

Brown Emergency Medicine

American Society of Regional Anesthesia and Pain Medicine (ASRA)
Reg Anesth Pain Med. 2018; 43(2):150-153

- Αλλεργία στη Lidocaine (αμίδιο): Αλλάζουμε ομάδα (εστέρες)

AMIDE GROUP (2 i´s)	ESTER GROUP (1 i)
Lidocaine	~~Cocaine~~
Mepivacaine	Procaine
Bupivacaine	Chloroprocaine
Etidocaine	Tetracaine
Prilocaine	Benzocaine

Νευροληπτοαναλγησία- καταστολή (sedation)

Intravenous sedation should be offered to patients, provided there are no contraindications. (Grade B)

Sedative drugs should be titrated to provide the desired depth of sedation. (Grade B)
The desired depth of sedation is one in which verbal contact is possible at all times. (Grade D)

Combination opioid and midazolam sedation should be considered in patients to improve bronchoscopic tolerance. (Grade B)

Short-acting agents (such as fentanyl or alfentanil) should be used to minimise post-procedural sedation. (Grade D)

When combination sedatives are used, opioids should be administered first and allowed time to become maximally effective before administration of any other agent. (Grade D)

Du Rand IA, et al. Thorax 2013;68:i1–i44

Drug	Dose Initial	Supplemental	Onset	Peak	Duration	Adverse efects	Antagonists
Fentanyl	25–50 µg	25 µg	3–5 min	5 min	1–2 h	Respiratory depression, nausea, vomiting	Naloxone
Alfentanil	250 µg	250 µg	Immediate	Immediate	1–2 h		
Morphine	2.5 m	2.5 m	5–10 min	15–30 mi	1–6 h		
Midazolam	2–2.5 mg	1 mg at 2–5 min interval	30–60 s	5–10 min	30–120 min	Respiratory depression, hypotension	Flumazenil 0.2 mg
Lorazepam	1.5–2 mg		8–15 min	15–30 min	8 h		
Diazepam	5–10 mg		1 min	2–3 min	1–3 h		
Propofol 1%	10–50 mg titrated to effect	25% of initial dose	30–60 s	2 min	4–8 min	Respiratory depression, bradycardia, hypotension, pain at the injection site	No antagonist available
Fospropofol	6.5 mg·kg–1	1.6 mg·kg–1	6.5 min	12 min	17 min		

Question I.12: What is the official and commonly accepted name for the epiglottis shown.

A. Omega or infantile epiglottis
B. Horseshoe epiglottis
C. U-shaped epiglottis
D. Normal appearing epiglottis

In children and in some adults, the epiglottis is elongated and curved.- omega epiglottis

http://www.bronchoscopy.org/education

Είσοδος βρογχοσκοπίου- Μύτη- Στόμα- Επιγλωττίδα- Λάρυγγας







- Έλεγχος σύγκλισης χορδών.
- Το γλωσσοφαρυγγικό και το πνευμονογαστρικό νεύρο νευρώνουν το λάρυγγα.
- Ο άνω λαρυγγικός κλάδος του πνευμονογαστρικού παρέχει την αισθητική νεύρωση του λάρυγγα.
- Ερεθισμός του μπορεί να οδηγήσει σε κλείσιμο των φωνητικών χορδών, για αυτό χρειάζεται τοπική αναισθησία.

Πολύποδες φωνητικών χορδών



Γαστροοισοφαγική παλινδρομήση



Ca φωνητικών χορδών





Παράλυση φωνητικής χορδής



Τραχεία- κυρία τρόπιδα

- Διάμετρος 16 με 20 mm
- 18–22 χόνδρινα ημικρίκια στο πρόσθιο μέρος.
- Το οπίσθιο μέρος είναι η υμενώδης μοίρα (λείες μυικές ίνες)
- Η κυρία τρόπιδα είναι στο επίπεδο του 5^{ου} θωρακικού σπονδύλου.΄Οξύαιχμη.

Τραχεία δίκην ξίφους (saber sheath)- ΧΑΠ



Τραχειακός βρόγχος- 0.1–3%



Αποπλάτυνση κυρίας τρόπιδας





Δυναμική σύγκλιση αεραγωγών- μείωση του αυλού κατά 50% κατά την εκπνοή (ΧΑΠ)



Τραχειομαλακία- αστάθεια των χόνδρων- κατάρρευση του πρόσθιου και οπισθίου τμήματος (συγγενής ή επίκτητη)



Μη επεμβατικός μηχανικός αερισμός σε τραχειομαλακία και e-DAC

- Καταστολή όπως στη συμβατική
- Έναρξη 20΄προ της βρογχοσκόπησης

Ξεκινάμε με:

- IPAP σε 14–15 cm και EPAP στα 5 cm σε bi-pressure systems
- 10 cm σε support pressure
- 5 cm όταν χρησιμοποιούμε CPAP



Τραχειο-οισοφαγικό συρίγγιο





- Έντονος βήχας με πυώδεις εκκρίσεις, ιδιαίτερα μετά από κατάποση
- Πνευμονία εξ εισροφήσεως
- Επιδείνωση της κλινικής κατάστασης του ασθενούς

Question II.14: The appearance of the airway stricture seen in the Figure should be described as

A. Simple
B. Hourglass
C. Complex



Στένωση τραχείας μετά από διασωλήνωση- Post intubation tracheal stenosis

- Γυναικα 45 ετών, αιμορραγικο ΑΕΕ- διασωλήνωση.
- Αποσωλήνωση- εισπνευστικός και εκπνευστικός συριγμός
- Αναπνευστική δυσχέρεια
- Τραχειοστομία



Παθογένεση της στένωσης τραχείας μετά από διασωλήνωση



>20-30mm Hg

Ισχαιμία βλεννογόνου

↓

Βλάβη στους χόνδρους

↓

Χονδρίτιδα → Ουλώδης ιστός

↘ Απώλεια στήριξης του χόνδρου



Ο αέρας στον αεροθάλαμο του τραχειοσωλήνα μπορεί να προκαλέσει έλκη και ουλώδη ιστό στο βλεννογόνο, οδηγώντας σε απλή ή σύνθετη στένωση

Δημιουργία ουλώδους ιστού περιμετρικά σαν ιστό- το τοίχωμα δεν καταστρέφεται



Μακρύτερη στένωση, με ανώμαλα όρια. Συμμετέχει το τραχειακό τοίχωμα και οι χόνδροι με έντονη ουλοποιήση
Συχνά παρατηρείται τραχειομαλακία



Σύνθετη στένωση με κοκκιώματα, παρουσία ράμματος από πρότερη επέμβαση

Virtual bronchoscopy





‘Ακαμπτη βρογχοσκόπηση

Τοποθέτηση stent τ. Dumon

Question IV.24: A thoracic surgery colleague has referred a patient for bronchoscopic removal of a foreign body located in RB2. On bronchoscopy, you will find the foreign body in

A. The posterior basal segment of the right lower lobe.
B. The medial basal segment of the right lower lobe.
C. The anterior segment of the left upper lobe.
D. The posterior segment of the right upper lobe.
E. The anterior segment of the right upper lobe.

Οι βρόγχοι αριθμούνται από 1-10
(1-3 άνω λοβός, 4-5 μέσος ή γλωσσίδα, 6-10 κάτω λοβός)
Οι τρόπιδες ονομάζονται C-1 or C-2.



http://www.bronchoscopy.org/education

- Ταξινόμηση των βρόγχων κατά Boyden.

Table 4.1. *Boyden surgical classification / Jackson–Huber classification*

Right bronchial tree		Left bronchial tree	
RUL		LUL	
B1	Apical	Upper division	
B2	Posterior	B1/2	Apicoposterior
B3	Anterior	B3	Anterior
RML		Lingular	
B4	Lateral	B4	Superior
B5	Medial	B5	Inferior
RLL		LLL	
B6	Superior	B6	Superior
B7	Medial basal	B7/8	Anteromedial
B8	Anterior basal	B9	Lateral basal
B9	Lateral basal	B10	Posterior basal
B10	Posterior		

Note: RUL, right upper lobe; LUL, left upper lobe; RML, right middle lobe; RLL, right lower lobe; LLL, left lower lobe.



YouTube

Search

Bronchoscopy Step by Step Walkthrough 1-6

https://www.youtube.com/watch?v=phRv73Ik7fI

http://www.bronchoscopy.org/education

A chest CT scan should be performed prior to a diagnostic bronchoscopy in patients with suspected lung cancer. (Grade D)



Du Rand IA, et al. Thorax 2013;68:i1–i44

Ο πήχης είναι ψηλά!

A diagnostic level of 85% should be attainable when definite endobronchial tumor is visible. (Grade B)

At least five biopsy samples should be taken when endobronchial tumor is visible to maximise diagnostic yield and the volume of biopsy tissue and to allow for tumor phenotyping and genotyping. (Grade D)

When endobronchial tumor is visible, brushings and washings can increase the diagnostic yield of the procedure. (Grade D)

Du Rand IA, et al. Thorax 2013;68:i1–i44

Κατηγοριοποίηση των βλαβών- βλεννογόνια/υποβλεννογόνια διήθηση- εξωφυτικοί όγκοι

Πλακώδες ca



Μικροκυτταρικό ca

- Εξωφυτική ανάπτυξη σε κεντρικό αεραγωγό- απόφραξη
- Νέκρωση
- Νεόπλαστα αγγεία (πιθανή αιμορραγική διάθεση στη βιοψία)
- Υποβλεννογόνια διήθηση



- Η **επιτυχής αιμόσταση** είναι πλέον απαραίτητη για την ασφαλή και επιτυχή έκβαση της βρογχοσκόπησης.

- **Σύγχρονες «θερμές» τεχνικές :**

1. **Ηλεκτροκαυτηρία (Electrocautery)**

2. **Argon Plasma Coagulation (APC)**

ERBE ICC 200
Electrosurgical Unit

Ηλεκτροκαυτηρία-μέθοδος επαφής



- Χρησιμοποιεί θερμότητα παραγόμενη από ηλεκτρισμό για να επιτευχθεί **αιμόσταση ή καταστροφή ιστού** ανάλογη με τη θερμική ενέργεια που θα αποδοθεί (συνήθως 25± 50 Watts)
- Βάθος διείσδυσης 0,2-1,9 mm.
- Χρησιμοποιείται με καθετήρα, λαβίδα ή θηλιά (snare) για μισχωτούς-πολυποειδείς όγκους

- Απόλυτη αντένδειξη είναι η παρουσία βηματοδότη (απαιτείται γνωμοδότηση του αρρυθμιολόγου) και η παρουσία εξωαυλικής βλάβης (πίεση εκ των έξω).

Argon Plasma Coagulation-μέθοδος μη επαφής

- Η θερμική ενέργεια μεταδίδεται μέσω ιονισμένου αερίου Αργόν το οποίο απελευθερώνεται σε μικρή απόσταση (non-contact mode) από τον βλεννογόνο.
- Εξαπλώνεται ομοιογενώς καθώς έλκεται από περιοχές με μεγαλύτερη περιεκτικότητα σε νερό (άρα και αίμα), και μικρότερη ηλεκτρική αντίσταση.
- Δεν σταματά σε καμπυλώσεις του βλεννογόνου ή σε πλάγιες επιφάνειες.

Υποβλεννογόνιες βλάβες-EBNA



- Η διαγνωστική ακρίβεια βιοψιών-ψήκτρας είναι μεγαλύτερη για τις εξωφυτικές βλάβες ενώ για τις υποβλεννογόνιες μπορεί να βελτιωθεί με τη χρήση **(E)-TBNA {**ευαισθησία κ.μ.ο. 80% (68-91%)**.**
- 55 ασθενείς - αύξηση της διαγνωστικής ευαισθησίας τους από το 76% στο 96% με την προσθήκη EBNA ενώ σε 20% ήταν η μόνη διαγνωστική εξέταση.



Dasgupta A. et al. Chest 1999;115:1237-1241

Question V.8 Moderate bleeding has occurred during an endobronchial biopsy of a nodule lying in the left main bronchus. The patient was immediately moved into the position shown. This is commonly called the

A. Fetal position
B. Trigger position
C. The safety position
D. The awkward position

Η πλάγια decubitus θέση με τον πάσχοντα πνεύμονα κάτω.
Το αίμα παραμένει κάτω με τη βαρύτητα.
Προστατεύεται η αντίθετη πλευρά.
Το αίμα αφαιρείται πιο εύκολα από τον φάρυγγα.

http://www.bronchoscopy.org/education

Μαζική αιμόπτυση

Παρά το γεγονός ότι η μαζική αιμόπτυση αποτελεί πιθανή θανατηφόρο κατάσταση, δεν υπάρχει σαφής συμφωνία στον ακριβη ορισμό.

Ποσότητα αίματος από 100 ml/24 h ως και 1,000 ml/24 h έχει προταθεί ως το όριο της μαζικής αιμόπτυσης.
Σημαντική αιμόπτυση αναφέρεται ποσότητα > 200 ml/24 h.

Περισσότερο σημαντικός είναι ορισμός που στηρίζεται σε σημαντικές κλινικές επιπτώσεις της αιμόπτυσης:

Απόφραξη του αεραγωγού και αιμοδυναμική αστάθεια
Αιμόπτυση που χρειάζεται μετάγγιση
Αιμόπτυση που χρήζει νοσηλείας
Αιμόπτυση που οδηγεί σε διασωλήνωση λόγω απόφραξης αεραγωγού.
Υποξυγοναιμία (Pa O2 < 60 mm Hg)
Θάνατος

Sakr I., Dutau H. Respiration 2010;80:35-58

Μαζική αιμόπτυση- αιτίες

Table 1. Major causes of massive hemoptysis

Infectious	Mycobacteria (particularly tuberculosis) Fungal infections (including mycetoma) Necrotizing pneumonia and lung abscess (*Klebsiella pneumoniae, Pseudomonas aeruginosa, Staphylococcus aureus, Streptococcus pneumoniae,* other *Streptococcus* spp. and *Actinomyces* spp.) Bacterial endocarditis with septic emboli Parasitic (paragonimiasis, hydatid cyst)
Neoplastic	Bronchogenic carcinoma Endobronchial tumors (carcinoid, adenoid cystic carcinoma) Pulmonary metastases Sarcoma
Pulmonary	Bronchiectasis (including cystic fibrosis) Chronic bronchitis Alveolar hemorrhage and underlying causes
Vascular	Pulmonary artery aneurysm (Rasmussen aneurysm, mycotic, arteritis) Bronchial artery aneurysm Pulmonary infarct (embolism) Pulmonary hypertension Congenital cardiac or pulmonary vascular malformations Airway-vascular fistula Arteriovenous malformations Mitral stenosis Left-ventricular failure
Vasculitis	Wegener's granulomatosis Goodpasture's syndrome Behçet's disease Systemic lupus erythematosus
Trauma	Induced by diagnostic bronchoscopy (brushing/biopsy) Related to interventional pulmonology procedures (dilation, metallic stent placement, high-dose brachytherapy) Catheter-induced PA rupture Blunt or penetrating chest injury Transtracheal procedures
Hemato-logical	Coagulopathy (congenital, acquired or iatrogenic) Platelet disorders
Drugs and toxins	Penicillamine Solvents Crack cocaine Trimellitic anhydride Bevacizumab
Miscella-neous	Endometriosis Lymphangioleiomatosis Broncholithiasis Cryptogenic Foreign body aspiration Lung transplantation

11-19% παραμένουν κρυπτογενείς

Sakr I., Dutau H. Respiration 2010;80:35-58

Διερεύνηση αιμόπτυσης- CT θώρακος

Η CT είναι ανώτερη διαγνωστικά από τη CXR, και συγκρίσιμη με τη βρογχοσκόπηση για την αναγνώριση του σημείου αιμορραγίας (σωστή εντόπιση σε 70–88.5%).

Η CT είναι πολύ περισσότερο αποτελεσματική από τη βρογχοσκόπηση για την εξακρίβωση του αιτίου αιμορραγίας (60–77 vs. 2.5–8%).



Diagnostic and Interventional Imaging (2015) 96, 775—788

Η βρογχοσκόπηση αναγνωρίζει το σημείο αιμορραγίας σε 73–93% των επεισοδίων σοβαρής αιμόπτυσης.

Η διαγνωστική αξία είναι χαμηλότερη σε ήπια ή μέτρια αιμόπτυση.



Καθετήρας με αεροθάλαμο και αποσπώμενη κεφαλή τύπου Freitag

Αφαίρεση θρόμβων με κρυόδιο

Το πρωτοξείδιο του αζώτου είναι το πιο χρησιμοποιούμενο-επιτυγχάνει θερμοκρασίες < -40 C. (Φαινόμενο Joule—Thomson)





DiBardino M et al. Annals ATS Volume 13 Number 8 August 2016

Ανίχνευση αγγειοδυσπλασίας-αιμαγγειώματος με HD video- Naroow Band Imaging και αντιμετώπιση με καυτηρίαση

Blue and green wavelengths are selected by optical filters, with the elimination of red light. These lights with narrowed bandwidths penetrate the superficial mucosal structures and are better absorbed by hemoglobin, providing an enhancement of mucosal features and blood vessels (capillaries from superficial mucosal layer, deeper mucosal and submucosal vessels)

Clin Endosc 2015;48:476-480

Question IV.14: Bronchoalveolar lavage is most helpful for providing histologic confirmation of diagnosis in all of the following diseases **except**

A. Chronic eosinophilic pneumonia
B. Histiocytosis X
C. Lymphangitic spread of carcinoma to the lungs
D. Pulmonary alveolar proteinosis
E. Invasive aspergillosis

BAL σε μη λοιμώδεις νόσους

Table 9.3. *Role of BAL in noninfectious diseases*

Diagnostic	Suggestive, not definitive
Pulmonary alveolar proteinosis	Alveolar hemorrhage
Langerhans cell histiocytosis	Asbestosis
Lipoid pneumonia/Fat embolism	Berylliosis
Malignancy	Drug-induced disease (amiodarone)
	Eosinophilic pneumonias
	Hypersensitivity pneumonitis
	Sarcoidosis
	Silicosis

Adapted from: *Am J Respir Crit Care Med.* 1990;142:481–486.



PAP- χρώση PAS
DAH- Σιδηροφάγα μακροφάγα (Hemosiderin-laden macrophages) και σταδιακή αιματηρή εμφάνιση του BAL.
Σαρκοείδωση: λόγος CD4-CD8 ≥4 έχει θετική προγνωστική αξία 94%.
Ιστιοκύττάρωση: Χρώση S-100 protein, OKT-6 (CD-1 antigen) μονοκλωνικό αντίσωμα.
Πνευμονίτιδα υπερευαισθησίας : μειωμένο CD4-CD8 <1 λόγω αύξησης suppressor T cells.

Introduction to Bronchoscopy. Armin Ernst. Cambridge Press 2009

BAL σε λοιμώδεις/μη λοιμώδεις νόσους

Table 9.4. *Disease associations with BAL studies/findings*

Microbiologic	Cytologic
KOH: fungal	Sulfur granules: actinomycetes
India ink: *Cryptococcus*	Hemosiderin-laden macrophages (alveolar hemorrhage) >20% red blood cells
Modified acid fast: *Nocardia*	Foamy macrophages: amiodarone, alveolar proteinosis, *Pneumocystis*
Ziehl-Neelsen, Auramine O: *Mycobacteria*	Malignancies
Silver methenamine: *Pneumocystis*/fungal	Oil Red O stain: fat embolism
Gram stain: predominant bacterial	Sudan stain: lipoid pneumonia
	>20% eosinophils: acute or chronic eosinophilic pneumonia, allergic bronchopulmonary aspergillosis, Churg–Strauss toxic inhalation, drug reaction
	Increased CD4-to-CD8 ratio: sarcoidosis, berylliosis, asbestosis, connective tissue disease
	Decreased CD4-to-CD8 ratio: hypersensitivity pneumonitis, silicosis, drug-induced disease, rheumatoid disease
	Periodic acid-Schiff: alveolar proteinosis
	Flow cytometry: lymphoma

Introduction to Bronchoscopy. Armin Ernst. Cambridge Press 2009

Διάγνωση λοίμωξης σε ανοσοεπαρκείς ασθενείς

Bronchoscopy may be considered in patients **with non-resolving or slowly resolving pneumonia**, especially if they are current or ex smokers and older than 50 years. (Grade C)

If bronchoscopy is performed for community-acquired pneumonia, BAL specimens should be sent for **legionella PCR and atypical pathogens**. (Grade C)

Bronchoscopy may be considered if the patient is suspected to have **TB** when sputum smear is negative. (Grade C)

In cases of suspected TB, **BAL, bronchial aspirates and post-bronchoscopy sputum** appear to be complementary and should all be analysed. (Grade C)



Du Rand IA, et al. Thorax 2013;68:i1–i44

Διάγνωση λοίμωξης σε ανοσοκατεσταλμένους

In patients with pulmonary infiltrates who are immunocompromised and in whom tuberculosis (TB) is considered unlikely, BAL alone is usually sufficient to achieve a diagnosis (Grade C)

In patients suspected to have invasive aspergillosis, BAL should be sent for microscopy for hyphae and fungal culture; a BAL galactomannan test should be considered to further improve diagnostic yield. (Grade C)
In invasive aspergillosis, BAL is reported to show characteristic hyphae in 34–64% and to be positive on cultures for Aspergillus in 23–85% of cases

TBLB and EBB for invasive aspergillosis may be avoided if BAL galactomannan test is available due to the high sensitivity and specificity of the latter and inherent risks with the biopsies. (Grade C)

Bronchoalveolar Lavage as diagnostic Modality for Pneumocystis jirovecii Pneumonia



Pneumocystis jirovecii cysts from bronchoalveolar lavage, stained with Toluidine blue stain

Το BAL έχει ευαισθησία μεταξύ 90% και 98% για P jiroveci.
Πρότερη χρήση εμπειρικής αντιμικροβιακής αγωγής μειώνει την ευαισθησία στο 64%.
Αμφοτερόπλευρη δειγματοληψία BAL έχει μεγαλύτερη διαγνωστική ακρίβεια συγκρινόμενη με την ετερόπλευρη λήψη.
BAL από τους άνω λοβούς έχει μεγαλύτερη ευαισθησία συγκριτικά με μέσο ή κάτω λοβό.



Courtesy of Dr Praveen Jha, Radiopaedia.org, rID: 17273

Golden, Jeffrey A et al. CHEST 90;1:18 - 22

Σταδιοποίηση NSCLC

Αφού έχει τεθεί η υποψία ή διάγνωση NSCLC πρέπει να εκτιμήσουμε την έκταση (στάδιο) της νόσου.

1. Υπάρχει απομακρυσμένη μετάσταση ή ο όγκος περιορίζεται στο θώρακα?
2. Αν περιορίζεται στο θώρακα, υπάρχει διήθηση των μεσοθωρακικών λεμφαδένων? Η διήθηση του μεσοθωρακίου θα επηρεάσει την στρατηγική θεραπείας.
3. Ασθενείς με στάδιο IA, IB, IIA, IIB μπορούν να ωφεληθούν από χειρουργική εκτομή του όγκου
4. Ασθενείς με στάδιο IIIA, IIB, IV συνήθως ωφελούνται από ΧΜΘ/ΑΚΘ.

TNM: Lung cancer

T/M	Subgroup	N0	N1	N2	N3
T1	T1a	Ia	IIa	IIIa	IIIb
	T1b	Ia	IIa	IIIa	IIIb
T2	T2a	Ib	IIa	IIIa	IIIb
	T2b	IIa	IIb	IIIa	IIIb
T3	$T3_{>7}$	IIb	IIIa	IIIa	IIIb
	$T3_{Inv}$	IIb	IIIa	IIIa	IIIb
	$T3_{Satell}$	IIb	IIIa	IIIa	IIIb
T4	$T4_{Inv}$	IIIa	IIIa	IIIb	IIIb
	$T4_{Ipsi\ Nod}$	IIIa	IIIa	IIIb	IIIb
M1	$M1a_{Contra\ Nod}$	IV	IV	IV	IV
	$M1a_{Pl\ Disem}$	IV	IV	IV	IV
	M1b	IV	IV	IV	IV

AJCC TNM Staging System, 7th Ed. (2010)

Προσδιορισμός του σταδίου (N):

- **N1:** Ομόπλευροι (της εξεργασίας) περιβρογχικοί, πυλαίοι και ενδοπνευμονικοί λεμφαδένες.
- **N2:** Ομόπλευροι μεσοθωρακικοί και/ή υποτροπιδικοί λεμφαδένες
- **N3:** -Ετερόπλευροι μεσοθωρακικοί και πυλαίοι
  -Ομόπλευροι ή ετερόπλευροι τραχηλικοί ή υπερκλείδιοι λεμφαδένες

- **Λεμφαδενική διόγκωση ορίζεται η αύξηση του μεγέθους ενός καλά διακριτού λεμφαδένα στον κοντό άξονα > 1 cm.**





Chest. 2013;143(5_suppl):e211S-e250S.
doi:10.1378/chest.12-2355

Η Σταδιοποίηση προβλέπει την επιβίωση



Prognosis according to the N category.

A: overall survival by cN category (any cT, cM0) using the IASLC classification. Reproduced from Rusch et al.[3]

B: overall survival by pN category (any pT, pM0) using the IASLC classification. Reproduced from Rusch et al.

[3] 1-yr = 1-year overall survival.

CHEST

Chest. 2009;136(1):260-271. doi:10.1378/chest.08-0978





Superior Mediastinal Nodes

- 1 Highest Mediastinal
- 2 Upper Paratracheal
- 3 Pre-vascular and Retrotracheal
- 4 Lower Paratracheal (including Azygos Nodes)

N_2 = single digit, ipsilateral
N_3 = single digit, contralateral or supraclavicular

Aortic Nodes

- 5 Subaortic (A-P window)
- 6 Para-aortic (ascending aorta or phrenic)

Inferior Mediastinal Nodes

- 7 Subcarinal
- 8 Paraesophageal (below carina)
- 9 Pulmonary Ligament

N_1 Nodes

- 10 Hilar
- 11 Interlobar
- 12 Lobar
- 13 Segmental
- 14 Subsegmental

Σταθμός 4R (δεξιοί κατώτεροι παρατραχειακοί)

ΑΝΩ ΟΡΙΟ	ΚΑΤΩ ΟΡΙΟ	ΜΕΣΗ ΓΡΑΜΜΗ
Κατώτερο σημείο επαφής της ανώνυμης (ή βραχιοκεφαλικής) φλέβας με την τραχεία	Κατώτερο όριο της αζύγου φλέβας	Αριστερό πλάγιο όριο της τραχείας
Με οδηγό σημείο την κυρία τρόπιδα στο βρογχικό δένδρο: TBNA-EBUS		





Δεξιά ανώνυμη φλέβα

Κάτω κοίλη φλέβα

Αορτικό τόξο



Σταθμός 4L (αριστεροί κατώτεροι παρατραχειακοί)

ΑΝΩ ΟΡΙΟ	ΚΑΤΩ ΟΡΙΟ	ΜΕΣΗ ΓΡΑΜΜΗ
Άνω όριο του αορτικού τόξου	Άνω όριο της αριστερής πνευμονικής αρτηρίας	Αριστερό πλάγιο όριο της τραχείας
Με οδηγό σημείο την κυρία τρόπιδα στο βρογχικό δένδρο: TBNA, EBUS, EUS		





Ανιούσα αορτή

Πνευμονική αρτηρία

Κατιούσα αορτή



Σταθμός 7 (υποτροπιδικοί)

ΑΝΩ ΟΡΙΟ	ΚΑΤΩ ΟΡΙΟ
Κυρία τρόπιδα	Κατώτερο όριο του διαμέσου βρόγχου δεξιά και ανώτερο όριο του κάτω λοβαίου βρόγχου αριστερά
Με οδηγό σημείο την κυρία τρόπιδα στο βρογχικό δένδρο: TBNA, EBUS, EUS	









Radiographic group A: patients with mediastinal infiltration so extensive that the discrete lymph nodes can no longer be discerned or measured

Radiographic group B: patients with measurable discrete mediastinal node enlargements.

Radiographic group C: the presence of a central tumor or suspected N1 disease makes the chance of N2,3 nodal involvement high (20%-25%) despite normal-sized nodes. Further confirmation is needed

Radiographic group D: patients with a peripheral clinical stage I tumor, where the chance of either distant metastases or mediastinal involvement is quite low (6-10% with regards to T1 or T2 stage)

Accuracy of CT scanning for staging of the mediastinum in patients with lung cancer (ACCP guidelines 2013).

First Author	Year	No.	Tech	Prev	Sens	Spec	PPV	NPV
Eggeling[137]	2002	73	CE	70	82	50	79	55
Wallace[143]	2001	121	CE	69	87	35	75	54
Marom[88]	1999	79	CE	56	59	86	84	63
Vansteenkiste[150]	1998	56	CE	50	86	79	80	85
Aaby[156]	1995	57	…	44	72	91	86	81
Schillaci[136]	2003	83	CE	42	69	75	67	77
Vansteenkiste[151]	1998	68	CE	41	75	63	58	78
Primack[157]	1994	159	CE	38	63	86	73	79
Turkmen[130]	2007	59	CE	36	43	66	41	68
Laudanski[141]	2001	92	CE	33	60	73	51	79
Yokoi1[58]	1994	113	CE	33	62	80	61	81
Gdeedo[153]	1997	100	CE	32	63	57	41	76
Bury[155]	1996	53	CE	32	71	81	63	85
McLoud[159]	1992	143	CE	31	64	62	44	79
Pieterman[90]	2000	102	CE	31	75	66	50	85
Yen[127]	2008	96	CE	31	47	80	52	77
Osada[146]	2001	335	CE	30	56	93	77	83
Jolly[160]	1991	336	CE	30	71	86	69	87
Subedi[47]	2009	91	CE	29	50	86	59	81
Buccheri[154]	1996	80	CE	28	64	74	48	84
Pozo-Rodriguez[24]	2004	132	CE	27	86	67	49	93
Kiernan[138]	2002	92	CE	27	64	94	80	88
Reed[19]	2003	302	CE	25	37	91	58	81
Nosotti[139]	2002	87	CE	25	64	88	64	88
Dunagan[144]	2001	72	CE	25	50	87	56	84
Kimura[135]	2003	203	CE	24	63	97	88	89
Yi1[29]	2007	143	CE	24	65	89	65	89
Suzuki[149]	1999	440	CE	23	33	92	56	82
Bury[152]	1997	64	CE	22	79	84	58	93
De Wever[44]	2007	50	CE	22	91	72	48	97
Webb[162]	1991	154	CE	21	52	69	31	84
Cole[161]	1993	150	…	21	26	81	26	81
Takamochi[132]	2005	71	CE	21	20	89	33	81
Kamiyoshihara[145]	2001	546	CE	20	33	90	46	84
Takamochi[147]	2000	401	CE	20	30	82	30	83
Lee[126]	2009	182	CE	20	36	79	30	83
Yang[128]	2008	122	CE	20	52	73	33	86
Kelly[134]	2004	69	CE	19	46	86	(43)[a]	87
Saunders[148]	1999	84	…	18	20	90	(30)[a]	84
Nomori[133]	2004	80	…	18	50	95	(70)[a]	90
Ebihara[131]	2006	205	CE	15	32	83	(26)[a]	87
Poncelet[142]	2001	62	CE	15	56	68	(23)[a]	90
Von Haag[140]	2002	52	CE	12	50	65	(16)[a]	91
Median: prevalence > 30					**67**	**74**	**62**	**79**
Median: prevalence 21-30					**63**	**87**	**58**	**84**
Median: prevalence ≤ 20					**41**	**83**	**30**	**87**
Summary: Median		**7,368**		**30**	**55**	**81**	**58**	**83**

- Sensitivity measures the proportion of positives that are correctly identified as such (e.g. the percentage of sick people who are correctly identified as having the condition).
- Specificity measures the proportion of negatives that are correctly identified as such (e.g. the percentage of healthy people who are correctly identified as not having the condition).

			Sens	Spec	PPV	NPV
Median: prevalence > 30			**67**	**74**	**62**	**79**
Median: prevalence 21-30			**63**	**87**	**58**	**84**
Median: prevalence ≤ 20			**41**	**83**	**30**	**87**
Summary: Median	**7,368**	**30**	**55**	**81**	**58**	**83**

Chest. 2013;143(5_suppl):e211S-e250S. doi:10.1378/chest.12-2355

Accuracy of integrated PET-CT scanning for staging of the mediastinum in patients with lung cancer (ACCP guidelines 2013).

First Author	Year	No.	Prev	Sens	Spec	PPV	NPV
Cerfolio[200]	2004	40	100	75	…	…	…
Plathow[204]	2008	52	73	100	100	100	100
Fischer[361]	2011	79	33	85	100	100	93
Lee[202]	2009	41	32	38	89	63	76
Yi[209]	2008	150	30	62	94	82	85
Maziak[29]	2009	167	29	48	93	74	82
Subedi[47]	2009	91	26	92	85	69	97
Yi[129]	2007	143	24	56	100	100	88
Carnochan[76]	2009	194	23	42	87	50	83
Lee[40]	2007	126	22	86	81	56	95
De Wever[44]	2007	50	22	73	82	53	91
Lee[126]	2009	182	20	81	73	42	94
Yang[128]	2008	122	20	52	73	33	86
Perigaud[203]	2009	51	20	40	85	40	85
Billé[199]	2009	159	19	48	93	(63)[a]	88
Toba[207]	2010	42	19	100	88	(67)[a]	100
Usuda[208]	2011	63	17	36	92	(50)[a]	87
Sanli[205]	2009	78	14	82	90	(56)[a]	97
Shin[206]	2008	184	13	48	95	(58)[a]	93
Summary: median		**2,014**	**22**	**62**	**90**	**63**	**90**

Chest. 2013;143(5_suppl):e211S-e250S. doi:10.1378/chest.12-2355

Πλεονεκτήματα vs. Μειονεκτήματα του PET CT

- Συνολικά 20% παραπάνω ασθενείς χαρακτηρίζονται σωστά ως έχοντες απομακρυσμένες (M1) ή λεμφαδενικές (N2,N3) μεταστάσεις συγκριτικά με την κλασσική σταδιοποίηση.
- Αν δεν επιβεβαιωθούν ιστολογικά, τα ευρήματα του PET CT μπορεί λανθασμένα να απομακρύνουν ασθενείς από πιθανή ριζική χειρουργική αντιμετώπιση.
- Στις κλινικές μελέτες με PET scan αυτό συνέβη σε ποσοστό 5-42%
- Το PET CT είναι λιγότερο ευαίσθητο σε βλάβες < 7-10 mm.

Η σύσταση της ACCP είναι:

3.1.2. In patients with an imaging finding (eg, by PET) suggestive of a metastasis, further evaluation of the abnormality with tissue sampling to pathologically confirm the clinical stage is recommended prior to choosing treatment (Grade 1B)**.**

Chest. 2013;143(5_suppl):e211S-e250S. doi:10.1378/chest.12-2355

Revised ESTS guidelines (2014) for preoperative mediastinal lymph node staging for non-small-cell lung cancer

- Addition of PET to staging adds to CT with overall sensitivity of 80–90% and specificity of 85–95%.
- PET–CT has a high NPV for detecting mediastinal nodal disease in peripherally located NSCLC.
- **Exceptions** include:
- 1. Suspected N1 nodes,
- 2. Tumor >3 cm
- 3. Centrally located tumor without suspected nodes on CT or PET scan.
- 4. Substantial difference in the rate of mediastinal nodal disease between adenocarcinoma and other tumour histology (risk ratio 2.72).
- 5. High 18F-Fluorodeoxyglucose (FDG) uptake in the primary lesion was associated with a greater risk of occult nodal metastasis.



For tumors >3 cm (mainly adenocarcinoma with high FDG uptake), or suspected N1, further mediastinal staging techniques providing histology should be considered.

European Journal of Cardio-Thoracic Surgery 45(2014)787–798

Πως θα ταυτοποιήσουμε ιστολογικά τα ευρήματα της CT και/ή PET CT? Μεσοθωρακοσκοπηση

First Author	Year	No.	Stage	Thoro	Prev	Sens	Spec[a]	PPV[a]	NPV
TM									
Hammoud[233]	1999	1,369	cN0-3	Sel	36	85	(100)[a]	(100)[a]	92
Lemaire[230]	2006	1,362	cN0-3	Sys	29	86	(100)[a]	(100)[a]	95
Coughlin[243]	1985	1,259	cN0-3	Sel	29	92	(100)[a]	(100)[a]	97
Luke[242, b]	1986	1,000	cN0-2	Sel	39	85	(100)[a]	(100)[a]	91
De Leyn[238]	1996	500	cN0-2	Sys	39	76	(100)[a]	(100)[a]	87
Anraku[227]	2010	352	cN0-3	Sys	37	92	(100)[a]	(100)[a]	96
Page[243, b]	1987	345	cN0-2	Sel	48	73	(100)[a]	(100)[a]	80
Dillemans[239]	1994	331	cN0-3	Sys	41	72	(100)[a]	(100)[a]	84
Brion[244]	1985	153	cN0-2	Sel	35	67	(100)[a]	(100)[a]	85
Fibla[231]	2006	142	cN0-3	Sel	42	67	(100)[a]	(100)[a]	80
Jolly[160]	1991	136	cN0-2	Sel	54	92	(100)[a]	(100)[a]	91
Ratto[125]	1990	123	cN0-2	Sel	33	88	(100)[a]	(100)[a]	94
Ebner[236, b]	1999	116	cN0-2	Sys	50	81	(100)[a]	(100)[a]	82
Annema[228]	2010	110	cN0-3	Sel	46	80	(100)[a]	(100)[a]	86
Gdeedo[155]	1997	100	cN0-3	Sys	32	78	(100)[a]	(100)[a]	91
Riordain[240]	1991	74	cN0-2	Sel	50	81	(100)[a]	(100)[a]	84
Aaby[156]	1995	57	cN0-3	Sys	44	84	(100)[a]	(100)[a]	89
Block[226]	2010	54	cN0-3	Sel	44	88	(100)[a]	(100)[a]	91
Kim[256]	2011	750	cN0	Sys	15	44	(100)[a]	(100)[a]	91
Choi[251]	2003	291	cN0	Sys	15	44	(100)[a]	(100)[a]	91
Meyers[229]	2006	169	cN0	Sel	8	38	(100)[a]	(100)[a]	95
Cerfolio[232]	2006	153	cN0-1	Sys	14	32	(100)[a]	(100)[a]	90
Deneffe[245]	1983	124	cN0	Sel	31	68	(100)[a]	(100)[a]	88
Park[225]	2010	78	cN0	Sys	8	50	(100)[a]	(100)[a]	96
Gurses[234]	2002	67	cN0	Sys	15	40	(100)[a]	(100)[a]	93
Leschber[222]	2008	52	cN0	Sys	19	…	(100)[a]	…	81
Median: cN0-3			***cN0-3***	*39% sys*	**40**	**83**			**90**
Median: cN0			***cN0***	*75% sys*	**16**	**47**			**91**
Median: sys			*46% cN0*	***sys***	**27**	**74**			**91**
Median: sel			*17% cN0*	***sel***	**39**	**81**			**91**
Summary TM: median		**9267**			**33**	**78**	**(100)[a]**	**(100)[a]**	**91**
VAM									
Venissac[223]	2003	154	cN2-3	Sys	71	97	(100)[a]	(100)[a]	94
Kimura[258]	2007	209	cN0-3	Sel	31	78	(100)[a]	(100)[a]	91
Lardinois[246]	2003	195	cN0-3	Sys	34	87	(100)[a]	(100)[a]	92
Kimura[135]	2003	125	cN0-3	Sys	36	85	(100)[a]	(100)[a]	92
Sayar[249]	2011	104	cN0-2	Sel	29	90	(100)[a]	(100)[a]	96
Anraku[227]	2010	89	cN0-3	Sys	22	95	(100)[a]	(100)[a]	99
Leschber[222]	2008	119	cN0	Sys	17	…	(100)[a]	…	83
Summary VAM: median		**995**			**31**	**89**	**(100)[a]**	**(100)[a]**	**92**
LA									
Zielinski[264]	2007	256	cN0-2	Compl	31	94	(100)[a]	(100)[a]	97
Witte[265]	2006	130	cN0-2	Compl		94	(100)[a]	(100)[a]	99
Summary ALL: median		**10,648**			**34**	**81**	**(100)[a]**	**(100)[a]**	**91**

- Χειρουργική αίθουσα-video μεσοθωρακοσκόπιο
- Γενική αναισθησία
- Νοσηρότητα 2% - Θνητότητα 0.08%
- Μη πρόσβαση σε σταθμούς 5, 6, 7p, 8,9

Sens	Spec	PPV	NPV
81	(100)[a]	(100)[a]	91

Διαβρογχική παρακέντηση (TBNA)

Lymph Node	Site
1. Anterior carina	First and second intercartilage interspace from lower trachea at about 12-1 o'clock position
2. Posterior carina	Posterior portion of carina at about 5-6 o'clock position
3. Right paratracheal	Second to fourth intercartilaginous interspace of lower trachea at about 1-2 o'clock position
4. Left paratracheal (aortic pulmonary window)	First or second intercartilaginous interspace from lower trachea at about 9 o'clock position
5. Right main bronchus	First or second intercartilaginous interspace from proximal right main bronchus at about 12 o'clock position
6. Left main bronchus	First or second intercartilaginous interspace from proximal left main bronchus at about 12 o'clock position
7. Right upper hilar	Anterior portion of right upper lobe spur
8. Subcarina	Medial wall of right main bronchus at about 9 o'clock position, proximal to level of right upper lobe orifice
9. Right lower hilar	Lateral or anterior wall of bronchus intermedius at about 3 o'clock position and 12 o'clock position near or at level of right middle lobe orifice
10. Subsubcarina	Medial wall of bronchus intermedius at about 9 o'clock position, proximal to level of right middle lobe orifice
11. Left hilar	Lateral wall of left lower lobe bronchus at about 9 o'clock, at level of superior segment orifice of left lower lobe





4R

7

Ko Pen Wang Chest. 1994;106(2):588-593.

Transbronchial needle aspiration of the mediastinum in patients with lung cancer.

Inclusion criteria: studies reporting test characteristics of TBNA of mediastinal nodes/tissue in patients with lung cancer, involving ≥ 20 patients from 1980 to 2011.

Study	Year	No.	Stage	Prev	Sens	Spec[a]	PPV[a]	NPV
Fernandez-Villar[277]	2010	280	cN1-3	…	68	(100)[a]	(100)[a]	10
Utz[294]	1993	61	cN0/N2	100	56	(100)[a]	(100)[a]	…
Selcuk[291]	2003	26	cN2-3	100	100	(100)[a]	(100)[a]	…
Wilsher[298]	1996	24	cN2-3	96	90	(100)[a]	(100)[a]	…
Katis[279]	1998	76	cN2-3	95	74	(100)[a]	(100)[a]	(20)[b]
Bilaceroglu[273]	1998	134	cN1-3	88	75	(100)[a]	(100)[a]	(36)[b]
Melloni[282]	2009	51	cN2-3	88	76	(100)[a]	(100)[a]	(33)[b]
Shannon[293]	1996	24	cN2-3	88	90	(100)[a]	(100)[a]	(60)[b]
Stratakos[294]	2008	77	cN1-3	86	88	(100)[a]	(100)[a]	(58)[b]
Schenk[288]	1993	64	cN2-3	86	56	(100)[a]	(100)[a]	(82)[b]
Wang[297]	1983	39	cN2-3	86	76	(100)[a]	(100)[a]	(71)[b]
Schenk[290]	1989	29	cN2-3	86	80	(100)[a]	(100)[a]	(44)[b]
Medford[281]	2010	79	cN2-3	84	79	(100)[a]	(100)[a]	(58)[b]
Mak[280]	2004	24	cN1-3	83	60	(100)[a]	(100)[a]	(33)[b]
Vansteenkiste[276]	1994	80	cN2	79	79	(100)[a]	(100)[a]	55
Rodriguez de castro[286]	1997	80	cN2-3	78	66	(100)[a]	(100)[a]	45
Harrow[278]	2000	264	cN1-3	72	93	99	99	80
Shah[292]	2006	129	cN1-3	71	68	(100)[a]	(100)[a]	56
Rodriguez de castro[275]	1995	56	cN2-3	70	77	(100)[a]	(100)[a]	70
Patelli[283]	2002	182	cN2	67	98	(100)[a]	(100)[a]	83
Rong[287]	1998	49	cN1-3	66	100	(100)[a]	(100)[a]	100
Rakha[284]	2010	182	cN1-3	57	84	(100)[a]	(100)[a]	70
Garpestad[274]	2001	32	cN1-3	57	83	(100)[a]	(100)[a]	67
Schenk[289]	1986	88	cN2-3	39	50	96	89	75
Bernasconi[176]	2006	113	cN2-3	31	54	(100)[a]	(100)[a]	91
Wallace[296]	2008	138	cN2-3	30	36	(100)[a]	(100)[a]	78
Ratto[285]	1988	47	cN2	30	14	(100)[a]	(100)[a]	73
Summary: median		**2,408**		**81**	**78**	**(100)[a]**	**(100)[a]**	**77**

Chest. 2013;143(5_suppl):e211S-e250S. doi:10.1378/chest.12-2355



- «Ιστολογική»(19-gauge) βελόνα για ιστό και κυτταρολογικά δείγματα.
- Καθημερινή χρήση της βελόνας ευαισθησία 88% και ειδικότητα 94% για NSCLC και SCLC

Stratakos G., et al. CHEST 2008; 133:131–136

- Αυτή η τεχνική μπορεί να εφαρμοστεί στην καθημερινή πρακτική με μεγάλη διαγνωστική ακρίβεια και ελάχιστες επιπλοκές (< 1%)
- Όμως, μόνο μικρός αριθμός βρογχοσκόπων την εφαρμόζει.
- Ο βασικός λόγος είναι η έλλειψη εκπαίδευσης και ο φόβος για παρακέντηση της βλάβης χωρίς την άμεση επισκόπηση (τυφλή τεχνική)

Διαβρογχική παρακέντηση (TBNA) με υπερηχολογική καθοδήγηση (EBUS-EUS)









Βελόνα 22Gauge

Η άγρια επιφάνεια επιτρέπει στη βελόνα να φαίνεται στον US





Προσαρμογή στο βρογχοσκόπιο

Δειγματοληψία

Πλακίδια και ιστικοί κύλινδροι



RAPID ON SITE EVALUATION (ROSE):

- Άμεση εκτίμηση του κυτταρολογικού υλικού της TBNA στην αίθουσα όπου πραγματοποιείται η βρογχοσκόπηση (real-time).
- Χρησιμοποιείται γρήγορη μέθοδος χρώσης 40 sec (συνήθως Diff Quik) για την εκτίμηση της επάρκειας του υλικού (αναγνώριση λεμφοκυττάρων, βρογχικού επιθηλίου ή παθολογικών κυττάρων).



Diff Quik smear, x 40 magnification, NSCLC

EBUS 6 LANDMARKS

SEARCH FOR THE LANDMARKS IN THIS ORDER:

4L → 7 → 10L → 10R → AZYGOS → 4R



N3→ N2 →N1

PFC 2015

Σταθμός 7

Σταθμός 10Rs

FIGURE 14. [Section 4.3.2.3] Endoscopic ultrasound-guided needle aspiration of the mediastinum in patients with lung cancer.

First Author	Year	No.	Stage	Thoro	Prev	Sens	Spec	PPV	NPV
Nadarajan[310]	2010	34	cN2-3	Sel	88	100	(100)[a]	(100)[a]	(100)[b]
Tournoy[312]	2008	100	cN0-3	Sys	83	95	(100)[a]	(100)[a]	(81)[b]
Wallace[143]	2001	121	cN2-3	Sel	79	87	(100)[a]	(100)[a]	68
Annema[323]	2004	36	cN2-3	Sys	78	93	(100)[a]	(100)[a]	80
Wiersema[305]	2001	29	cN2-3	Sel	76	100	(100)[a]	(100)[a]	100
Fritscher-Ravens[301]	2000	35	cN2-3	Lim	74	96	(100)[a]	(100)[a]	90
Annema[316]	2005	215	cN0-3	Sys	71	91	(100)[a]	(100)[a]	74
Larsen[324]	2002	29	cN2-3	Lim	69	90	(100)[a]	(100)[a]	82
Annema[306]	2010	551	cN2-3	Sys	66	83	(100)[a]	(100)[a]	75
Caddy[320]	2005	36	cN0-3	Sel	65	92	(100)[a]	(100)[a]	83
Kalade[311]	2008	33	cN1-3	Sel	64	95	(100)[a]	(100)[a]	92
Silvestri[304]	1996	26	cN0-3	Sys	62	89	(100)[a]	(100)[a]	82
Gress[302]	1998	24	cN2-3	Sel	58	93	(100)[a]	(100)[a]	90
Herth[307]	2010	139	cN1-3	Sel	52	89	(100)[a]	(100)[a]	82
Talebian[309]	2010	152	cN2-3	Sys	49	74	(100)[a]	(100)[a]	73
Sawhney[313]	2006	65	cN2-3	Sys	48	87	(100)[a]	(100)[a]	90
Fritscher-Ravens[179]	2003	33	cN0-3	Sys	48	88	(100)[a]	(100)[a]	89
Larsen[324]	2005	58	cN0-3	Sys	47	87	(100)[a]	(100)[a]	87
Eloubeidi[315]	2005	104	cN2-3	Sys	41	93	(100)[a]	(100)[a]	96
Eloubeidi[318]	2005	35	cN2-3	Sys	37	91	(100)[a]	(100)[a]	97
Annema[317]	2005	100	cN2-3	Sys	36	71	90	86	85
Wallace[322]	2004	69	cN0	Sys	36	61	(100)[a]	(100)[a]	82
LeBlanc[321]	2005	67	cN0	Sel	33	45	(100)[a]	(100)[a]	79
Wallace[296]	2008	138	cN2-3	Sys	30	69	(100)[a]	(100)[a]	88
Szlubowski[308]	2010	120	cN0	Sel	22	50	99	93	87
Fernandez-Esparrach[314]	2006	47	cN0	Sys	21	50	(100)[a]	(100)[a]	88
Median: prevalence > 80		% sys	50%			**96**			**90**
Median: prevalence 60-79		% sys	40%			**92**			**82**
Median: prevalence 40-59		% sys	71%			**88**			**88**
Median: prevalence 20-39		% sys	71%			**61**			**87**
Median: cN1-3		% sys	47%			**92**			**89**
Median: cN0-1		% sys	50%			**50**			**85**
Summary: median		**2,443**			**58**	**89**	**(100)[a]**	**(100)[a]**	**86**

Chest. 2013;143(5_suppl):e211S-e250S.
doi:10.1378/chest.12-2355

Προτείνεται EBUS TBNA σε PET (-) ασθενείς?

- 100 ασθενείς με αρνητικό PET και υψηλή υποψία για NSCLC.
- 156 λεμφαδενες παρακεντήθηκαν με διάμετρο 5-10 mm.
- 8 ασθενείς είχαν θετικούς λεμφαδένες για NSCLC:

-N0 σε N3 (1 ασθενής),
-N0 σε N2 (5 ασθενείς)
-N0 σε N1 (2 ασθενείς).

- 98.9% NPV

Herth et al CHEST 2008;133:887-891

- 113 ασθενείς με NSCLC N0 σε PET CT.
- EBUS-TBNA προεγχειρητικά.
- N2 Νόσος σε 17.6% (20 από 113).

Shingyoji et al, Annals of Thoracic Surgery 2014;98: 1762-1767

Αντιστοιχία θέσης βρογχοσκοπίου στο βρόγχο με US εικόνα και CT



BRONCHOSCOPY INTERNATIONAL™

www. Bronchoscopy.org

Άλλοι περιορισμοί EBUS TBNA

- Υψηλό κόστος εγκατάστασης/αναλωσίμων/ζημίας
- Ακατάλληλο για βρογχοσκόπηση ρουτίνας
- Δυσανεξία ασθενούς στην εξέταση-μερικές φορές είναι απαραίτητη διασωλήνωση πάρα τη χρήση μιδαζολάμης-φεντανύλης ή προποφόλης.
- Οι λεμφαδενικοί σταθμοί 5,6,8,9 δεν προσεγγίζονται (σταθμοί 8,9 προσεγγίζονται με το **EUS-FNA- υπερηχογαστροσκόπιο**).



Μέση ευαισθησία και ειδικότητα από κέντρα τα οποία πραγματοποίησαν έλεγχο με EBUS TBNA και EUS FNA

FIGURE 16. [Section 4.3.2.5] Real-time EBUS-TBNA and EUS-NA of the mediastinum in patients with lung cancer.

First Author	Year	No.	Stage	Thoro	Prev	Sens	Spec[a]	PPV[a]	NPV
Vilmann[353]	2005	31	cN0-3	sys	65	100	(100)[a]	(100)[a]	100
Annema[228]	2010	123	cN1-3	sys	54	82	(100)[a]	(100)[a]	80
Herth[307]	2010	139	cN1-3	sel	52	96	(100)[a]	(100)[a]	96
Ohnishi[352]	2011	110	cN0-3	sys	35	72	(100)[a]	(100)[a]	87
Hwangbo[335]	2010	150	cN2-3	sys	31	91	(100)[a]	(100)[a]	96
Wallace[296]	2008	138	cN2-3	sys	30	93	(100)[a]	(100)[a]	97
Szlubowski[308]	2010	120	cN0	sel	22	68	98	91	91
Median: prevalence 40-65						**96**			**96**
Median: prevalence 20-39						**82**			**94**
Summary: median		**811**			**33**	**91**	**(100)[a]**	**(100)[a]**	**96**

Chest. 2013;143(5_suppl):e211S-e250S. doi:10.1378/chest.12-2355

Συνδυαστική προσπέλαση μεσοθωρακίου από το EBUS βρογχοσκόπιο μέσω οισοφάγου και τραχείας

Combined Endoscopic-Endobronchial Ultrasound-Guided Fine-Needle Aspiration of Mediastinal Lymph Nodes Through a Single Bronchoscope in 150 Patients With Suspected Lung Cancer

Felix J. F. Herth, Mark Krasnik, Nicolas Kahn, Ralf Eberhardt and Armin Ernst

Chest 2010;138;790-794; Prepublished online February 12, 2010;
DOI 10.1378/chest.09-2149

The combined approach had a sensitivity of 96% and a negative predictive value of 95%, values higher than either approach alone. No complications occurred.
Conclusions: The two procedures can easily be performed with a dedicated linear endobronchial ultrasound bronchoscope in one setting and by one operator. They are complementary and provide better diagnostic accuracy than either one alone. The combination may be able to replace more invasive methods as a primary staging method for patients with lung cancer.

CHEST 2010; 138(4):790–794

Prof. J. T. Annema AMC, Amsterdam, Netherlands

ERS School on IP, Sotiria, Athens, September 25, 2014



Επανασταδιοποίηση με EBUS TBNA?

Endobronchial Ultrasound With Transbronchial Needle Aspiration for Restaging the Mediastinum in Lung Cancer

- 124 ασθενείς (IIIA) επανασταδιοποίηση με EBUS TBNA
- CT: 58(47%) stable, 66(53%) downstaged
- EBUS: Persistent 89(72%) nodal metastasis
- From 35 patients in whom no metastases were assessed by EBUS TBNA 28 had residual N2 disease
- Negative predictive value: 20%

Table 2. Outcomes for Restaging EBUS-TBNA According to Tumor Response After Chemotherapy As Assessed by RECIST Criteria

	Tumor Response Outcome (%)	
Criteria	Partial Response (n = 66)	Stable Disease (n = 58)
Sensitivity	77	75
Specificity	100	100
PPV	100	100
NPV	22	18
Accuracy	79	76

Abbreviations: EBUS-TBNA, endobronchial ultrasound–guided transbronchial needle aspiration; RECIST, Response Evaluation Criteria in Solid Tumors Group; PPV, positive predictive value; NPV, negative predictive value.

Λόγω της χαμηλής αρνητικής προγνωστικής αξίας προτείνεται μεσοθωρακοσκόπηση για επιβεβαίωση των αρνητικών EBUS TBNA

Herth et al J Clin Oncol 26:3346-3350

ANNALS OF THE AMERICAN THORACIC SOCIETY
A journal for clinicians, clinical researchers, and medical educators

Adequacy of Samples Obtained by EBUS-TBNA for Molecular Analysis in Patients with Non-Small Cell Lung Cancer: Systematic Review and Meta-Analysis

Gonzalo Labarca [1,2,3], Erik Folch[3,4], Michael Jantz[3,5], Hiren J. Mehta[3,5], Adnan Majid[3,6], Sebastian Fernandez-Bussy[3,7]

Table 6. Summary of findings: Molecular analysis from NSCLC using samples obtained by EBUS-TBNA.

Outcomes	Explanation, measure effect (95% CI)	Number of participants (studies)	Certainty of the evidence (GRADE)
EGFR (overall)	Samples achieved by EBUS-TBNA are 94.8% (93% to 96.4) adequate for EGFR analysis	2497 (28 observational studies)	⊕◯◯◯[1,2,3] VERY LOW
EGFR (Asia)	Samples achieved by EBUS-TBNA is Asia are 98.9% (98.1% to 99.7%) adequate for EGFR analysis	626 (8 observational studies)	⊕⊕◯◯[1,3] LOW
ALK (overall)	Samples achieved by EBUS-TBNA are 94.9% (89.4% to 98.8) adequate for ALK analysis	607 (10 observational studies)	⊕◯◯◯[1,2,3] VERY LOW

1: Risk of bias; 2: Imprecision; 3: Indirectness.

Conclusion: EBUS-TBNA has a high yield for molecular analysis of both EGFR and ALK mutations.

Περιφερικός (μονήρης) όζος πνεύμονα – SPN, Solitary Pulmonary Nodule:

Στρογγύλη ή ωοειδής πνευμονική σκίαση με δ ≤ 3cm (που περιβάλλεται από υγιές πνευμονικό παρέγχυμα).



- 27.3% pts (σύνολο 53.454) που συμμετείχαν στη National Lung Screening Trial (NLST) είχαν τουλάχιστον έναν όζο στην ΥΤ

NEJM. March 2015

Δεν είναι όλα κακοήθειες…

Adenocarcinoma

Chondroamartoma

Διαθωρακική παρακέντηση υπο CT καθοδήγηση

- Έχει διαγνωστικό εύρος 72-90%
- Μεγαλύτερη επίπτωση επιπλοκών από τις βρογχοσκοπικές μεθόδους.
- Προκαλεί Πνευμοθώρακα σε ποσοστό 30% και 5-10% αυτών θα χρειαστούν σωλήνα θωρακικής παροχέτευσης

Διαβρογχική παρακέντηση (TBB) για την προσπέλαση περιφερικών όζων-μαζών

- Η TBB έχει την μεγαλύτερη ευαισθησία 57% (21 μελέτες) ακολουθούμενη από το brushing 54% και το BAL 43%.
- Η TBNA σε μικρό αριθμό εργασιών αναφέρεται με ευαισθησία 65%.



Initial Diagnosis of Lung Cancer: ACCP Evidence-Based Clinical Practice guidelines)
M. Patricia Rivera and Atul C. Mehta Chest 2007;132;131-148



Βρογχοσκόπηση – TBB με ακτινοσκόπηση

All Methods:		< 2 cm LESION				> 2 cm LESION			
First Author	Year	N	Pos	Neg	Sens	N	Pos	Neg	Sens
Gasparini[110]	1995	195	82	113	**42**	300	169	131	**56**
Hattori[76]	1971	17	13	4	**76**	182	150	32	**82**
Baaklini[88]	2000	16	4	12	**25**	135	93	42	**69**
Wallace[122]	1982	65	3	62	**5**	78	24	54	**31**
Bandoh[130]	2003	25	8	17	**32**	72	50	22	**69**
Radke[106]	1979	21	6	15	**29**	76	49	27	**64**
Naidich[121]	1988	15	4	11	**27**	46	26	20	**57**
Trkanjec[129]	2003	17	9	8	**53**	33	27	6	**82**
McDougall[105]	1981	9	1	8	**11**	36	21	15	**58**
Stringfield[107]	1977	3	1	2	**33**	26	13	13	**50**
Summary		383	131	252	**34**	984	622	362	**63**

CHEST 2013; 143(5):e142S–e165S



Σχέση όγκου/βρόγχου κατά Tsuboi

- Η διαβρογχική βιοψία TBLB έχει μικρότερη διαγνωστική ακρίβεια σε βλάβες τύπου III και IV (0.25 vs 0.6).
- Σε ασθενείς που η λαβίδα ή η ψήκτρα δεν μπορούν να φθάσουν στο κέντρο του όγκου η TBNA προσφέρει αποκλειστική διάγνωση με ευαισθησία της τάξης του 80%.
- Σήμερα είναι δεκτό ότι σε βλάβες τύπου III και IV η TBNA είναι μέθοδος εκλογής

- ***Bilaceroglu et al. Respiration 1998;65:49-55***
- ***Shure D et al. Am Rev Respir Dis 1983;128:1090-1092.***

EBUS-radial

Table 1. Studies evaluating the diagnostic yield of radial probe endobronchial ultrasound since 2014

Study	Year	Pro/Retro*	Number of lesions	Diameter mean	Diameter median	Diagnostic yield %	Guide sheath	Fluoro*	Other tools*
Boonsarngsuk *et al.* [26]	2014	Retro	174	25.1 mm (SD 10.7 mm)	NR	79.9% overall	+	+	Forceps, brush, BAL
Ikezawa *et al.* [27]	2014	Retro	67	21 mm (SD 8 mm)	NR	57% overall	+	+	Double-hinged curette, forceps, brush
Chen *et al.* [28]	2014	Retro	467	NR	NR	69% overall	+/−	−	Forceps, brush, TBNA
Hayama *et al.* [29]	2015	Retro	965	NR	25 mm (range 6–107)	64.4% overall, 77.1% for malignancy	+	+	Forceps, brush, TBNA, VNB, ROSE
Durakovic *et al.* [30]	2015	Retro	147	28 mm (SD 18 mm)	NR	26.5% for malignancy	−	−	Forceps, brush, BW
Casutt *et al.* [31]	2015	Retro	51	25.8 mm (range 8–67 mm)	NR	72.5% overall	−	+	Forceps, brush, BAL/BW, TBNA
Chavez *et al.* [32]	2015	Retro	212	20.45 mm (SD 5.45 mm)	NR	67.5% for malignancy	+	+	Forceps, brush, TBNA, ROSE

*Pro, prospective; Retro, retrospective.
BAL, bronchoalveolar lavage; BW, bronchial washing; Fluoro, fluoroscopy; GS, guide sheath; NR, not reported; r-EBUS, radial probe endobronchial ultrasound; ROSE, rapid on-site evaluation; TBNA, transbronchial needle aspiration; VNB, virtual navigational bronchoscopy.

Variability 26,5-79,9% (εξαρτάται από το μέγεθος, περισσότερο όμως από τη θέση της βλάβης σε σχέση με το βρόγχο – σημαντικό το σημείο αεροβρογχογράμματος)

Curr Opin Pulm Med 2016, 22:000–000


Type I
Bronchus


Type II


Type III


Tumor

Διαβρογχική βιοψία με την καθοδήγηση radial EBUS

Πλεονεκτήματα

- Αύξηση του διαγνωστικού εύρους
- Μη απαραίτητη η ακτινοσκόπηση
- Χαμηλή πιθανότητα επιπλοκών

Μειονεκτήματα

- Η καθοδήγηση είναι περιορισμένη
- Επιπλέον κόστος (probe, guide sheath)
- Το εύρος περιορίζεται όταν SPN < 2cm

- *Stratakos et al, Annals of Thoracic Medicine 2017*
- *Schwarz Y et al. Chest 2006,*
- *Gildea TR et al. Am J Respir Crit Care Med 2006,*
- *Eberhardt R et al. Chest 2007,*
- *Makris D et al. Eur Respir J 2007,*
- *Eberhardt R et al. Am J Respir Crit Care Med 2007*

Non electromagnetic navigation - Lungpoint®



Procedure planning



Real-time synchronization

Non real-time virtual bronchoscopy navigation - Synapse®



Βρογχοσκοπική βιοψία μονήρων πνευμονικών όζων (SPN) χωρίς ενδοβρογχική διαδρομή.

BTPNA (Bronchoscopic transparenchymal needle aspiration) με το σύστημα Archimedes© (Broncus Medical).



1. Συγχρονισμός της βρογχοσκοπικής εικόνας με τη virtual bronchoscopy-εικονικό μονοπάτι



2. Δημιουργία Tunnel και διαστολή με αεροθάλαμο

3. Βιοψία με βελόνη υπό ακτινοσκοπική καθοδήγηση

Ion by Intuitive
A new robotic endoluminal platform* for minimally invasive peripheral lung biopsy

Athens welcomes

6th European Congress for Bronchology and Interventional Pulmonology

ECBIP 2021 6TH

Organized by

Hellenic Training Association for Interventional Pulmonology

EABIP

National & Kapodistrian University of Athens

endorsed by

HELLENIC THORACIC SOCIETY

ERS EUROPEAN RESPIRATORY SOCIETY

ECBIP

April 22- 24

2021

BACK UP SLIDES

Διάμεσες πνευμονοπάθειες- νεότερα δεδομένα (διαβρογχική κρυοβιοψία)

Διάγνωση διάμεσων πνευμονοπαθειων-IPF



An Official ATS/ERS/JRS/ALAT Clinical Practice Guideline: Treatment of Idiopathic Pulmonary Fibrosis: Executive Summary

An Update of the 2011 Clinical Practice Guideline

Ganesh Raghu, Bram Rochwerg, Yuan Zhang, Carlos A. Cuello Garcia, Arata Azuma, Juergen Behr, Jan L. Brozek, Harold R. Collard, William Cunningham*, Sakae Homma, Takeshi Johkoh, Fernando J. Martinez, Jeffrey Myers, Shandra L. Protzko, Luca Richeldi, David Rind, Moisés Selman, Arthur Theodore, Athol U. Wells, Henk Hoogsteden, and Holger J. Schünemann; on behalf of the ATS, ERS, JRS, and ALAT

American Journal of Respiratory and Critical Care Medicine Volume 192 Number 2| July 15 2015

A Surgical Lung Biopsy Showed a Pattern of Usual Interstitial Pneumonia





In-Hospital Mortality after Surgical Lung Biopsy for Interstitial Lung Disease in the United States

2000 to 2011

John P. Hutchinson, Andrew W. Fogarty, Tricia M. McKeever, and Richard B. Hubbard

Division of Epidemiology and Public Health, School of Medicine, University of Nottingham, Nottingham, United Kingdom

Table 2. In-Hospital Mortality after Surgical Lung Biopsy for Interstitial Lung Disease, by Year

Year of Biopsy	Total Admissions [*Deaths (% Mortality)*]	Elective Admissions [*Deaths (% Mortality)*]	Nonelective Admissions [*Deaths (% Mortality)*]
2000	822 (7.6)	120 (1.2)	591 (16.6)
2001	962 (7.5)	173 (2.2)	690 (17.6)
2002	923 (6.9)	196 (2.1)	727 (17.5)
2003	934 (7.3)	173 (2.1)	761 (17.6)
2004	875 (7.0)	131 (1.6)	745 (18.8)
2005	876 (6.6)	148 (1.7)	727 (16.6)
2006	876 (6.6)	139 (1.6)	736 (16.5)
2007	696 (5.4)	116 (1.3)	580 (14.1)
2008	715 (5.8)	135 (1.6)	580 (15.0)
2009	709 (5.5)	123 (1.5)	586 (13.2)
2010	696 (5.7)	139 (1.7)	557 (14.3)
2011	617 (4.9)	101 (1.2)	516 (13.5)
Total	9,700 (6.4)	1,695 (1.7)	7,796 (16.0)

- Around 12,000 surgical lung biopsies performed annually for interstitial lung disease in US.
- Two-thirds of which were performed electively.
- In-hospital mortality was 1.7% for elective procedures but significantly higher for non-elective procedures (16.0%).
- Male sex, increasing age, increasing comorbidity, open surgery, and a provisional diagnosis of idiopathic pulmonary fibrosis or connective tissue disease-related interstitial lung disease were risk factors for increased mortality.

Am J Respir Crit Care Med Vol 193, Iss 10, pp 1161–1167, May 15, 2016

Cryobiopsy Equipment:

1.Console and gas cylinder
2.Transfer line (connects the console to the probe)
3.Cryoprobe (flexible :1.9 mm/2.4 mm and rigid: 3.0 mm)
4.Pedal

ERBECRYO_CA

ERBECRYO_2



Efficacy and safety of cryobiopsy versus forceps biopsy for interstitial lung diseases and lung tumours: A systematic review and meta-analysis

OORMILA GANGANAH, Shu Liang GUO, Manu CHINIAH AND Yi Shi LI

Table 1 Characteristics of included studies

First author	Year	Number of subjects	Disease involved	Type of study
Babiak	2009	41—all subjects, FB and CB	ILDs	Retrospective
Schumann C.	2010	First 55—all subjects, FB and CB	Lung tumours	Randomized cohort
Aktas A.	2010	41—all subjects, CB and FB	Lung tumours	Prospective clinical trial
Griff	2011	33—15, CB; 18, FB	ILDs	Prospective clinical Trial
Hetzel J.	2012	563—282, CB; 281, FB	Lung tumours	Randomized single blinded multicentre
Schuhmann M.	2013	31—all subjects, CB and FB	Peripheral solitaryLung tumours	Randomized clinical trial
Chou C.L.	2013	75—all subjects, CB and FB	ILDs	Retrospective
Pajares	2014	77	ILDs	Randomized trial

CB, cryobiopsy; FB, forceps biopsy; ILDs, interstitial lung diseases.

Respirology (2016)

Efficacy and safety of cryobiopsy versus forceps biopsy for interstitial lung diseases and lung tumours: A systematic review and meta-analysis

OORMILA GANGANAH, Shu Liang GUO, Manu CHINIAH AND Yi Shi LI

Table 1 Characteristics of included studies

First author	Year	Number of subjects	Disease involved	Type of study	Specimen size	
					CB	FB
Babiak	2009	41—all subjects, FB and CB	ILDs	Retrospective	15.11 mm^2 (2.15–54.15 mm^2)	5.82 mm^2 (0.58–20.88 mm^2)
Schumann C.	2010	First 55—all subjects, FB and CB	Lung tumours	Randomized cohort	10.4 mm^2	5.2 mm^2
Aktas A.	2010	41—all subjects, CB and FB	Lung tumours	Prospective clinical trial	0.8 cm (0.3–4.0 cm)	0.2 cm (0.1–1.0 cm)
Griff	2011	33—15, CB; 18, FB	ILDs	Prospective clinical Trial	17.1 ± 10.7 mm^2	3.8 ± 4.0 mm^2
Hetzel J.	2012	563—282, CB; 281, FB	Lung tumours	Randomized single blinded multicentre		
Schuhmann M.	2013	31—all subjects, CB and FB	Peripheral solitaryLung tumours	Randomized clinical trial	11.17 mm^2	4.69 mm^2
Chou C.L.	2013	75—all subjects, CB and FB	ILDs	Retrospective	13.8 ± 3.8 mm	1.9 ± 0.6 mm
Pajares	2014	77	ILDs	Randomized trial	14.7 ± 11 mm^2	3.3 ± 4.1 mm^2

CB, cryobiopsy; FB, forceps biopsy; ILDs, interstitial lung diseases.



x **2-3**

Respirology (2016)

Efficacy and safety of cryobiopsy versus forceps biopsy for interstitial lung diseases and lung tumours: A systematic review and meta-analysis

OORMILA GANGANAH, Shu Liang GUO, Manu CHINIAH AND Yi Shi LI

Table 1 Characteristics of included studies

First author	Year	Number of subjects	Disease involved	Type of study	Specimen size		Diagnostic rate (%)	
					CB	FB	CB	FB
Babiak	2009	41—all subjects, FB and CB	ILDs	Retrospective	15.11 mm^2 (2.15–54.15 mm^2)	5.82 mm^2 (0.58–20.88 mm^2)	95.12	> 58.53
Schumann C.	2010	First 55—all subjects, FB and CB	Lung tumours	Randomized cohort	10.4 mm^2	5.2 mm^2	89.1	> 65.5
Aktas A.	2010	41—all subjects, CB and FB	Lung tumours	Prospective clinical trial	0.8 cm (0.3–4.0 cm)	0.2 cm (0.1–1.0 cm)	92.7	> 78
Griff	2011	33—15, CB; 18, FB	ILDs	Prospective clinical Trial	17.1 ± 10.7 mm^2	3.8 ± 4.0 mm^2		
Hetzel J.	2012	563—282, CB; 281, FB	Lung tumours	Randomized single blinded multicentre			95	> 85.1
Schuhmann M.	2013	31—all subjects, CB and FB	Peripheral solitaryLung tumours	Randomized clinical trial	11.17 mm^2	4.69 mm^2	61.2	> 48.4
Chou C.L.	2013	75—all subjects, CB and FB	ILDs	Retrospective	13.8 ± 3.8 mm	1.9 ± 0.6 mm	100	> 69.3
Pajares	2014	77	ILDs	Randomized trial	14.7 ± 11 mm^2	3.3 ± 4.1 mm^2	74.4	> 34.1

CB, cryobiopsy; FB, forceps biopsy; ILDs, interstitial lung diseases.

Respirology (2016)

DIAGNOSTIC YIELD

Study or Subgroup	cryobiopsy Events	cryobiopsy Total	forceps biopsy Events	forceps biopsy Total	Weight	Risk Ratio M-H, Random, 95% CI
1.2.2 Interstitial lung diseases						
Babiak et al	39	41	24	41	13.5%	1.63 [1.24, 2.12]
Pajares et al	29	39	13	38	7.3%	2.17 [1.35, 3.51]
Subtotal (95% CI)		**80**		**79**	**20.8%**	**1.77 [1.34, 2.32]**
Total events	68		37			
Heterogeneity: Tau² = 0.01; Chi² = 1.22, df = 1 (P = 0.27); I² = 18%						
Test for overall effect: Z = 4.08 (P < 0.0001)						
1.2.3 lung tumours						
Aktas et al	38	41	32	41	16.8%	1.19 [0.99, 1.43]
ChunLiang et al	75	75	52	75	18.1%	1.44 [1.24, 1.67]
Hetzel et al	268	282	239	281	21.0%	1.12 [1.06, 1.18]
Schuhmann M et al	19	31	15	31	7.7%	1.27 [0.80, 2.00]
Schumann et al	49	55	36	55	15.6%	1.36 [1.10, 1.68]
Subtotal (95% CI)		**484**		**483**	**79.2%**	**1.25 [1.10, 1.43]**
Total events	449		374			
Heterogeneity: Tau² = 0.01; Chi² = 12.99, df = 4 (P = 0.01); I² = 69%						
Test for overall effect: Z = 3.32 (P = 0.0009)						



Respirology (2016)

Transbronchial Cryobiopsy in Diffuse Parenchymal Lung Disease: Need for Procedural Standardization

Venerino Poletti[a, b] Jürgen Hetzel[c]

Table 1. Transbronchial cryobiopsy for diffuse lung disease: different approaches

First author [Ref.]	Year	OT	RB	LM	NI	GA+JV	GA/DS	LA	Bronchial blocker	Cryoprobe size, mm	Freezing time, s
Babiak [10]	2009	x					x		N	2.4	4
Pajares [11]	2010	x					x		N	2.4	3
Griff [12]	2011		x		x		x	x			
Kropski [13]	2013	x					x			1.9	4
Yarmus [14]	2013		x (10)	x (11)		x	x		Y	1.8	3
Fruchter [15]	2013				x			x	N	2.4	4
Fruchter [16]	2013				x			x	N	2.4	4
Fruchter [17]	2014				x			x	N	2.4	4
Casoni [18]	2014		x				x		Y	2.4	5/6
Pajares [19]	2014	x					x		Y	2.4	3/4
Poletti [7]	2014		x				x		Y	2.4	5/6
Griff [20]	2014		x		x		x	x	N	1.9	3/5
Gershman [22]	2015				x			x	N	2.4	4
Hagmeyer [23]	2015	x	x			x			N	2.4	4/5
Hernández-González [21]	2015	x					x		Y	1.9	3/4

OT = Orotracheal tube; RB = rigid bronchoscope; LM = laryngeal mask; NI = no intubation; GA = general anesthesia; JV = jet ventilation; DS = deep sedation; LA = local anesthesia; Y = yes; N = no; x = method used.

Respiration 2015;90:275–278

Table 2. Comparison between different transbronchial cryobiopsy modalities

	Intubation with OT	Intubation with RT	LM	NI
Need of GA/DS	Y	Y	Y	N
Control of bleeding	Y	Y	Y/N	N
Size of specimens	+++	+++	+	+
Size of OF	++	+++	+	–
Comfort for patients	+++	+++	+++	–
Technical effort	++	+++	+	–

OT = Orotracheal tube; RT = rigid tracheoscope; LM = laryngeal mask; NI = no intubation; GA = general anesthesia; DS = deep sedation; OF = operating field; Y = yes; N = no.

Respiration 2015;90:275–278